

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**Thermography in systemic sclerosis patients and other rheumatic diseases: Diagnosis, disease activity assessment, and therapeutic monitoring**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1723440> since 2020-01-16T15:45:43Z

*Published version:*

DOI:10.1016/j.autrev.2019.102449

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

**This is the author's final version of the contribution published as:**

[Autoimmun Rev.](#) 2019 Dec 12:102449. doi: 10.1016/j.autrev.2019.102449. [Epub ahead of print]

## **Thermography in systemic sclerosis patients and other rheumatic diseases: Diagnosis, disease activity assessment, and therapeutic monitoring.**

Sciascia S, Cecchi I, Massara C, Rossi D, Radin M, Ladehesa PL, Guiñazu F, Rubini E, Foddai SG, Alba P, Escudero A, Menegatti E, Roccatello D.

**The publisher's version is available at:**

<https://www.sciencedirect.com/science/article/pii/S1568997219302629?via%3Dihub>

**When citing, please refer to the published version.**

**Link to this full text:**

<http://hdl.handle.net/2318/1723440>

This full text was downloaded from iris-AperTO: <https://iris.unito.it/>

iris-AperTO

University of Turin's Institutional Research Information System and Open Access Institutional Repository

# **Thermography in systemic sclerosis patients and other rheumatic diseases: diagnosis, disease activity assessment, and therapeutic monitoring**

Sciascia S<sup>\*1</sup>, Cecchi I<sup>\*1,2</sup>, Massara C<sup>1</sup>, Rossi D<sup>1</sup>, Radin M<sup>1,2</sup>, Ladehesa Pineda L<sup>3</sup>, Guiñazu F<sup>4</sup>, Rubini E<sup>1,2</sup>, Foddai S<sup>1,2</sup>, Alba P<sup>4</sup>, Escudero A<sup>3</sup>, Menegatti E<sup>1,2</sup>, and Roccatello D<sup>1</sup>

**\*equally contributed**

1. Center of Research of Immunopathology and Rare Diseases- Coordinating Center of Piemonte and Valle d'Aosta Network for Rare Diseases, Department of Clinical and Biological Sciences, and SCU Nephrology and Dialysis, S. Giovanni Bosco Hospital, Turin, Italy
2. School of Specialization of Clinical Pathology, Department of Clinical and Biological Sciences, University of Turin, Italy
3. Department of Rheumatology, Reina Sofía University Hospital, Maimonides Biomedical Research Institute of Córdoba (IMIBIC), University of Córdoba, Córdoba, Spain
4. Rheumatology, Hospital of Córdoba, Córdoba, Argentina

## **Corresponding Author:**

Dario Roccatello, MD

Center of Research of Immunopathology and Rare Diseases - Coordinating Center of Piemonte and Valle d'Aosta Network for Rare Diseases, and SCU Nephrology and Dialysis, S. Giovanni Bosco Hospital, Piazza del Donatore di Sangue 3, 10154, Turin, Italy.

Email [dario.roccatello@unito.it](mailto:dario.roccatello@unito.it) - Tel +390112402051 Fax +390112402052

**Key words:** Systemic sclerosis, Raynaud's Phenomenon, Thermography, Autoimmune Disease, Rheumatic Disease, Imaging

Dear Sir,

The recent advancement of knowledge and management strategies of patients with autoimmune conditions such as systemic sclerosis (SSc), inflammatory arthritis, and systemic lupus erythematosus (SLE) achieved during the last years lead to the clear-cut improvement of patients' quality of life and reduced long-term disability [1–5]. Nowadays, new imaging techniques in autoimmune-rheumatic diseases are expanding the possibilities to achieve an early diagnosis, an appropriate follow-up, evaluation of disease activity, monitoring, and response to treatment in the affected patients. Among these techniques, infrared thermography represents a promising tool in order to assess vascular and joint involvement in autoimmune patients.

Since the initial enthusiasm in medical use of thermal photography over 2 decades ago, only few clinical practical applications were developed until past years, when the technical quality of available imaging systems advanced immensely. Improved data quality enhances the capability for detecting diagnostically relevant information in the thermal images and reduces human errors and variations in the image interpretation.

Modern thermal imaging systems are equipped to detect this radiation and can be used to form a dynamic map of temperature distribution at a surface. In brief, infrared camera generates images with thermal data in real time, and it is possible in a fraction of a second to image large areas of the human body, for example all the joints of both hands.

Thermal cameras images can be taken as a pseudo-measure of blood flow due to the convection of heat from the blood vessels in the skin and muscle in response to thermoregulation.

Moreover, as infrared thermography can give an indirect measure of blood flow and articular inflammation, it can be useful to assess peripheral vasculopathy and synovitis in active rheumatic arthritis, both at baseline and in response to treatment.

## 1. Arthritis

Synovitis represents the cardinal sign of arthritis, and therefore its assessment in patients with inflammatory arthritis is a major issue for proper treatment. The complementary methods of images

have become increasingly relevant, being the magnetic resonance imaging (MRI) and ultrasound (US) widely used for this purpose. Nevertheless, both techniques have several limitations and the need for an ideal imaging method is still a matter of research.

In this context, thermography represents a promising imaging tool which has been not yet exhaustively explored.

Among others, Collins and Cosh and Ring and Collins [6,7] showed that the surface temperature of an arthritic joint was related to the temperature in intra-articular joint, and to other serological markers of inflammation. More recently, Brenner and colleagues demonstrated that changes observed in the thermal signature correlated with disease severity in rodent models of mono-arthritis and systemic autoimmune arthritis, suggesting that the novel thermal imaging techniques could provide useful and objective measurements of joint inflammation based on two cardinal inflammatory signs: joint swelling/edema (joint diameters) and warmth [8] (Figure 1a). New thermal imaging devices have shown to be effective at detecting changes in temperature in the context of inflammation and arthritis in *in vivo* models [9].

An early detection of articular inflammation, even before a definitive diagnosis is established, can be crucial, since starting an appropriate treatment at early stages of the pathological process seems to prevent disease persistence, joint destruction, and long-term disability [10].

Assessment of thermal imaging could be an easy to use and cost-effective technique in the general clinical practice especially when firstly assessing a new onset of arthralgia in patients suspected for rheumatic diseases. Critically, it might represent a tool to investigate the presence of active joint inflammation in the armamentarium of general practitioners when considering referring a patient for a rheumatologic evaluation and when high resolution ultrasound examination is not available or pending. Nevertheless, larger studies are still needed to confirm and validate the usefulness of thermography in this setting as the previous studies centered mainly in exploring the thermographic differences between arthritic patients and healthy subjects.

## 2. Critical Digital Ischemia

Critical digital ischemia is a rare but serious complication of rheumatic diseases, mainly systemic sclerosis (SSc). It is characterized by the obstruction of the arterial flow, which causes severe pain at rest, compromising tissue viability, potentially leading to ulceration, gangrene, and ultimately amputation. In patients with SSc, this complication is the result of both progressive microangiopathy and digital artery vasculopathy [11], rather than to large vessel disorder which occurs in patients with atherosclerotic peripheral disease.

It is well recognized that microvascular changes can be easily demonstrated by nailfold capillaroscopy, which cannot provide yet functional measure of blood flow [12].

Clinical assessment and investigation need to center on establishing the underlying cause, in order to optimize treatment strategies. However, to date, the overall management of patients affected by critical digital ischemia is difficult for the treating clinician and new tools both for diagnosis and monitoring ongoing treatment are highly needed. Infrared thermal system has been previously applied as non-invasive monitoring tool that provides real-time screening information of tissue perfusion based on infrared thermal signals. Therefore, the use of thermal imaging might be an additional tool to assess critical digital ischemia and to monitor it during therapy (Figure 1b). Therefore, this approach might potentially provide helpful supplementary information about tissue perfusion in patients suffering from SSc and in those patients who experience this severe complication. Further studies are warranted to validate the diagnostic and the reproducibility of the approach.

### 3. Systemic Lupus Erythematosus

Systemic lupus erythematosus (SLE) is characterized by a broad spectrum of clinical manifestations enfolding different tissues and organs at the onset and during the course of the disease. Among these, the presence of synovitis, meaning “non-erosive arthritis” involving  $\geq 2$  peripheral joints, characterized by tenderness, swelling or effusion or “inflammatory synovitis” in  $\geq 2$  joints characterized by: a) swelling or effusion, or b) tenderness and 30 minutes of morning stiffness [13] are included in the classification criteria. The joint involvement in SLE is non-erosive, non-

deforming arthralgias/arthritis primarily affecting small joints of both hands, wrists and knees, and few patients show a major synovitis indistinguishable from rheumatoid arthritis (RA), the *so-called* “rhupus”.

A number of techniques are available in order to objectify inflammation and joint swelling, such as MRI, which however is expensive and not practical for repetitive use. The use of high-resolution US examination represents a reliable technique when performed by expert and trained operators. However, it is not always available outside rheumatologic practise or might request a further referral. Assessment of thermal imaging could be a useful, easy to perform and cost-effective tool to be applied in both general and rheumatic settings in order to improve the management of patients with SLE complaining for arthralgia, providing a fast and reproducible methods with the potential to guide therapeutic choices (Figure 1c).

#### 4. Systemic Sclerosis

SSc is a chronic autoimmune disorder with high morbidity and mortality. The pathogenesis of the disease is characterized by the presence of both micro- and macro-vascular damage and fibrosis. Skin lesions represent the leading feature of the disease and Raynaud’s phenomenon (RP) is the most frequent and early clinical manifestation. Digital vasculopathy severely affects SSc patients, negatively impacting on their daily functions. In addition, recent evidences have shown that the presence of digital ulcers at the beginning of the disease is associated with poor cardiovascular prognosis and decreased survival rate [14]. Nevertheless, adequate treatment strategies for the management of RP and skin manifestations are lacking, mainly due to the absence of reliable outcome measures in clinical trials [15]. Therefore, identifying new tools for an early diagnosis and monitoring of vasculopathy in SSc patients represents an urgent clinical need. Moreover, the recent development of new therapeutic agents makes this need even more important in order to assess their efficacy in preventing and treating skin and vascular involvement in this setting. In fact, as stated before, even if nailfold capillaroscopy has been already validated as an accountable tool for the

analysis of those structural changes that can differentiate between primary and secondary RP, it cannot provide functional analysis of blood flow.

Previous studies have evaluated the possible role of infrared thermography imaging for the assessment of vascular damage and RP in affected patients [16–19], but its role needs further validation in SSc setting and its use in monitoring the clinical evolution and/or response to therapy has been only anecdotally explored.

Therefore, in order to evaluate the efficacy of infrared thermography in the assessment of peripheral vasculopathy in a cohort of SSc patients treated with cyclic intravenous infusions with synthetic analogue of prostacyclin (Iloprost), we designed this pilot prospective study. Critically, taking into account the lack of standard protocol timing the use of synthetic prostanoids for SSc patients, we also aimed to identify those patients who might benefit from an intensified infusional treatment protocol with prostanoids. For the sake of this study, we enrolled patients diagnosed with SSc [20], attending the San Giovanni Bosco Hospital of Turin (Italy) for their routinely 28-days apart intravenous therapy with Iloprost based on the presence of severe secondary RP and/or digital ulcers. Thermographic evaluation of both hands was performed at baseline (T0) before prostanoid infusion, and at days 14 and 28 after the first prostanoid infusion (named T1 and T2, respectively). For the thermographic recordings, a digital infrared camera FLIR ONE (FLIR Systems Inc., USA) was used. The acquired thermal images were analyzed with software dedicated to thermographic analysis FLIR Tools (FLIR Systems Inc., USA). Camera was placed on horizontal surface at 1 m distance (field of view  $25^{\circ} \times 19^{\circ}$ ) perpendicular to skin surface. The significance of baseline differences was determined by the chi-squared test, Fisher's exact test or the unpaired t-test, as appropriate. A two-sided P-value  $<0.05$  was statistically significant. All statistical analyses were performed using SPSS version 19.0 (IBM, Armonk, NY, USA). As result, among the 26 enrolled patients, 24 (92.3%) were female, mean age was  $65.3 \pm 12.6$  years, and mean disease duration at time of data collection was  $12.4 \pm 6.6$  years. Fifteen patients (57.7%) presented with a limited



cutaneous form of SSc, while 11 patients (42.3%) showed a diffuse cutaneous form of SSc with a high rate of interstitial lung disease as high as 63.6%. The thermographic assessment showed a substantial stability of the temperature values when comparing T0 and T1 (mean differences of the right hands  $0.4 \pm 5.6$ ; mean differences of the left hands  $1.2 \pm 4.5$ ), while they are significantly reduced when comparing T1 and T2 (mean differences of the right hands  $-3.1 \pm 9.3$ ,  $p=0.049$ ; mean differences of the left hands  $-3.4 \pm 8.5$ ,  $p=0.012$ ) (Figure 2a). When stratifying according to clinical manifestation, a higher difference in temperature variations were observed between T1 and T2 in patients with diffuse cutaneous SSc, when compared to those with limited cutaneous SSc (mean of the differences of the right hands  $-5.0 \pm 11$ ; mean of left-hands differences  $-4.9 \pm 11.5$  Vs. mean right-hands differences  $-2.5 \pm 11$ ; mean left-hands differences  $-3 \pm 8.6$ ;  $p=0.035$  respectively) (Figure 2b).

## 5. Conclusion and future prospective

Autoimmune rheumatic diseases are a heterogeneous group of chronic pathologic conditions affecting joints and connective tissue, often leading to permanent damage and disability, lowering the quality of life of the affected patients. An early diagnosis, an appropriate therapeutic strategy and monitoring are therefore mandatory and highly needed. In this setting, infrared thermography might represent a reliable, non-invasive, portable, and cost-effective diagnostic tool both in expert centers and in general practice. Moreover, although limited, the results of this prospective pilot study seem very promising to reinforce the role of thermography as suitable outcome measure of novel treatments for SSc-related RP and vasculopathy in clinical trials. Moreover, these data show that patients with diffuse cutaneous form of SSc could benefit more from an intensified infusion protocol with prostanoids compared to those SSc patients with a limited cutaneous form of the disease. Indeed, further longitudinal studies are needed in order to assess the validity and the reproducibility of this technique.

**Conflict of interest:** None

**Funding:** None

## **Bibliography**

- [1] Yue X, Yu X, Petersen F, Riemekasten G. Recent advances in mouse models for systemic sclerosis. *Autoimmun Rev* 2018;17:1225–34. doi:10.1016/j.autrev.2018.06.013.
- [2] Sanz Pérez I, Martínez Valle F, Guillén-del-Castillo A, Roque Pérez A, Cuéllar Calàbria H, Pizzi MN, et al. Subclinical cardiovascular disease and Systemic Sclerosis: A comparison between risk charts, quantification of coronary calcium and carotid ultrasonography. *Autoimmun Rev* 2018;17:900–5. doi:10.1016/j.autrev.2018.03.015.
- [3] Cutolo M, Vanhaecke A, Ruaro B, Deschepper E, Ickinger C, Melsens K, et al. Is laser speckle contrast analysis (LASCA) the new kid on the block in systemic sclerosis? A systematic literature review and pilot study to evaluate reliability of LASCA to measure peripheral blood perfusion in scleroderma patients. *Autoimmun Rev* 2018;17:775–80. doi:10.1016/j.autrev.2018.01.023.
- [4] Rossi D, Russo A, Manna E, Binello G, Baldovino S, Sciascia S, et al. The role of nail-videocapillaroscopy in early diagnosis of scleroderma. *Autoimmun Rev* 2013;12. doi:10.1016/j.autrev.2012.11.006.
- [5] Giuggioli D, Manfredi A, Lumetti F, Colaci M, Ferri C. Scleroderma skin ulcers definition, classification and treatment strategies our experience and review of the literature. *Autoimmun Rev* 2018;17:155–64. doi:10.1016/j.autrev.2017.11.020.
- [6] Collins AJ, Ring EF, Cosh JA, Bacon PA. Quantitation of thermography in arthritis using multi-isothermal analysis. I. The thermographic index. *Ann Rheum Dis* 1974;33:113–5.
- [7] Collins AJ, Cosh JA. Temperature and biochemical studies of joint inflammation. A

preliminary investigation. *Ann Rheum Dis* 1970;29:386–92.

- [8] Brenner M. Thermal signature analysis as a novel method for evaluating inflammatory arthritis activity. *Ann Rheum Dis* 2006;65:306–11. doi:10.1136/ard.2004.035246.
- [9] Herrick AL, Murray A. The role of capillaroscopy and thermography in the assessment and management of Raynaud's phenomenon. *Autoimmun Rev* 2018;17:465–72. doi:10.1016/j.autrev.2017.11.036.
- [10] Wevers-de Boer KVC, Heimans L, Huizinga TWJ, Allaart CF. Drug therapy in undifferentiated arthritis: a systematic literature review. *Ann Rheum Dis* 2013;72:1436–44. doi:10.1136/annrheumdis-2012-203165.
- [11] Herrick AL. The pathogenesis, diagnosis and treatment of Raynaud phenomenon. *Nat Rev Rheumatol* 2012;8:469–79. doi:10.1038/nrrheum.2012.96.
- [12] Cutolo M, Sulli A, Smith V. Assessing microvascular changes in systemic sclerosis diagnosis and management. *Nat Rev Rheumatol* 2010;6:578–87. doi:10.1038/nrrheum.2010.104.
- [13] Petri M, Orbai A-M, Alarcón GS, Gordon C, Merrill JT, Fortin PR, et al. Derivation and validation of the Systemic Lupus International Collaborating Clinics classification criteria for systemic lupus erythematosus. *Arthritis Rheum* 2012;64:2677–86. doi:10.1002/art.34473.
- [14] Mihai C, Landewé R, van der Heijde D, Walker UA, Constantin PI, Gherghe AM, et al. Digital ulcers predict a worse disease course in patients with systemic sclerosis. *Ann Rheum Dis* 2016;75:681–6. doi:10.1136/annrheumdis-2014-205897.
- [15] García de la Peña Lefebvre P, Nishishinya MB, Pereda CA, Loza E, Sifuentes Giraldo WA, Román Ivorra JA, et al. Efficacy of Raynaud's phenomenon and digital ulcer pharmacological treatment in systemic sclerosis patients: a systematic literature review. *Rheumatol Int* 2015;35:1447–59. doi:10.1007/s00296-015-3241-1.

- [16] Wilkinson JD, Leggett SA, Marjanovic EJ, Moore TL, Allen J, Anderson ME, et al. A Multicenter Study of the Validity and Reliability of Responses to Hand Cold Challenge as Measured by Laser Speckle Contrast Imaging and Thermography: Outcome Measures for Systemic Sclerosis-Related Raynaud's Phenomenon. *Arthritis Rheumatol* (Hoboken, NJ) 2018;70:903–11. doi:10.1002/art.40457.
- [17] Merkel PA, Herlyn K, Martin RW, Anderson JJ, Mayes MD, Bell P, et al. Measuring disease activity and functional status in patients with scleroderma and Raynaud's phenomenon. *Arthritis Rheum* 2002;46:2410–20. doi:10.1002/art.10486.
- [18] Pauling JD, Shipley JA, Harris ND, McHugh NJ. Use of infrared thermography as an endpoint in therapeutic trials of Raynaud's phenomenon and systemic sclerosis. *Clin Exp Rheumatol* n.d.;30:S103-15.
- [19] Allen J, Howell K. Microvascular imaging: techniques and opportunities for clinical physiological measurements. *Physiol Meas* 2014;35:R91–141. doi:10.1088/0967-3334/35/7/R91.
- [20] van den Hoogen F, Khanna D, Fransen J, Johnson SR, Baron M, Tyndall A, et al. 2013 Classification Criteria for Systemic Sclerosis: An American College of Rheumatology/European League Against Rheumatism Collaborative Initiative. *Arthritis Rheum* 2013;65:2737–47. doi:10.1002/art.38098.